

I claim:

1. An improved method for treating male erectile dysfunction comprising:

administering to a male patient an effective amount of the combination of sildenafil citrate and papaverine hydrochloride, said papaverine hydrochloride administered enterally.

2. The treatment method of claim 1 wherein:

sildenafil citrate is administered in dosages of between approximately 25 and approximately 100 milligrams approximately every four to six hours, and

papaverine hydrochloride is administered enterally in dosages of between approximately 50 and approximately 450 milligrams per day.

3. The treatment method of claim 2 including:

the contemporaneous administration of between approximately 200 milligrams and approximately 1,200 milligrams of pentoxifylline per day.

4. The treatment method of claim 2 wherein:

approximately 100 milligrams of sildenafil citrate is administered.

5. The treatment method of claim 2 wherein:

approximately 150 milligrams of papaverine hydrochloride is administered twice a day.

1 6. The treatment method of claim 4 wherein:

2 approximately 150 milligrams of papavarine hydrochloride is administered twice a day.

1 7. The treatment method of claim 1 further comprising:

2 an effective amount of pentoxifylline.

1 8. The treatment method of claim 7 wherein

2 between approximately 200 milligrams and approximately 1,200 milligrams per day of  
3 pentoxifylline is administered.

1 9. The treatment method of claim 8 wherein:

2 approximately 400 milligrams of pentoxifylline is administered per day.

1 10. The treatment method of claim 4 including:

2 the contemporaneous administration of approximately 400 milligrams of pentoxifylline  
3 three times a day.

1 11. The treatment method of claim 5 including:

2 the contemporaneous administration of approximately 400 milligrams of pentoxifylline  
3 three times a day.

1 12. The treatment method of claim 6 including:

2 the contemporaneous administration of between approximately 200 milligrams and  
3 approximately 1,200 milligrams of pentoxifylline per day.

1 13. The treatment method of claim 12 wherein:

2 approximately 400 milligrams of pentoxifylline is administered three times a day.

1 14. The treatment method of claim 1 further comprising:

2 an effective amount of zinc monomethionine aspartate.

1 15. The treatment method of claim 14 wherein:

2 between approximately 25 and approximately 75 milligrams per day of zinc  
3 monomethionine aspartate is administered per day.

1 16. The treatment method of claim 15 wherein:

2 approximately 25 milligrams per day of zinc monomethionine aspartate is administered.

1 17. The treatment method of claim 4 wherein:

2 approximately 25 milligrams per day of zinc monomethionine aspartate is administered.

1 18. The treatment method of claim 5 wherein:

2 approximately 25 milligrams per day of zinc monomethionine aspartate is administered.

1 19. The treatment method of claim 6 wherein:

2 approximately 25 milligrams per day of zinc monomethionine aspartate is administered.

1 20. The treatment method of claim 10 wherein:

2 approximately 25 milligrams per day of zinc monomethionine aspartate is administered.

1 21. The treatment method of claim 11 wherein:

2 approximately 25 milligrams per day of zinc monomethionine aspartate is administered.

1 22. The treatment method of claim 12 wherein:

2 approximately 25 milligrams per day of zinc monomethionine aspartate is administered.

1 23. The treatment method of claim 13 wherein:

2 approximately 25 milligrams per day of zinc monomethionine aspartate is administered.

1 24. An improved method for treating male erectile dysfunction comprising:

2 administering to a male patient an effective amount of the combination of sildenafil citrate,  
3 papavarine hydrochloride, pentoxifylline and zinc monomethionine aspartate.

1 25. An improved method for treating male erectile dysfunction comprising:

2 administering to a male patient the combination of

3 sildenafil citrate in dosages of between 25 and approximately 100 milligrams

4 during a period of between approximately four and approximately six hours,  
5 papavarine hydrochloride, administered enterally, in dosages of between  
6 approximately 50 and approximately 450 milligrams per day, and  
7 between approximately 200 milligrams and approximately 1,200 milligrams of  
8 pentoxifylline per day.

1 26. The treatment method of claim 25 including:

2 the contemporaneous administration of between approximately 25 and approximately 75  
3 milligrams per day of zinc monomethionine aspartate.

1 27. An improved method for treating male erectile dysfunction comprising:

2 administering to a male patient the combination of  
3 sildenafil citrate in dosages of between 25 milligrams and approximately 100  
4 milligrams approximately every four to six hours,  
5 papavarine hydrochloride, administered enterally, in dosages of between  
6 approximately 50 milligrams and approximately 450 milligrams per day,  
7 pentoxifylline in dosages of between approximately 200 milligrams and  
8 approximately 1,200 milligrams per day, and  
9 zinc monomethionine aspartate in dosages of between approximately 25  
10 milligrams and approximately 75 milligrams per day.